

RECEIVED
CENTRAL FAX CENTER

FEB 16 2007

Amendments to the Claims

1. (currently amended) A method of releasing a protein of interest from host cells comprising
 - a. suspending the host cells in a buffer, and
 - b. adding a detergent to the suspended cells wherein the detergent is an amphipathic charged amine or an amphipathic charged amine oxide, and wherein the detergent is in a concentrated form of at least 3 times the desired final concentration of the detergent after dilution in the buffer of step a.
2. (currently amended) A method of releasing a protein of interest from host cells comprising
 - a. suspending the host cells in a buffer, and
 - b. adding a detergent to the suspended cells, wherein the detergent is selected from the group consisting of : tributylphosphate, dimethyldodecylamine, dimethyltridecylamine, dimethylundecylamine, dimethyldidecylamine, dimethyltetradecylamine, dimethylhexadecylamine, dimethyldecylamineoxide, dimethylundecylamineoxide, dimethyldidecylamineoxide, dimethyltetradecylamineoxide and dimethyltridecylamineoxide, and wherein the detergent is in a concentrated form of at least 3 times the desired final concentration of the detergent after dilution in the buffer of step a.
3. (currently amended) The method of claim 1 or 2, wherein the detergent is not dimethyltridecylamine.
4. (currently amended) The method of claim 3, further comprising the step of adding glycerol to the suspended host cells, and wherein the glycerol is in a concentrated form of at least 3 times the desired final concentration of the glycerol after dilution in the buffer of step 1.
5. (original) The method of claim 4, wherein the detergent comprises a final concentration of between 0.1 to 10 percent.

6. (original) The method of claim 4, wherein the glycerol comprises a final concentration of between 0.6 to 15 percent.
7. (original) The method of claim 5, wherein the glycerol comprises a final concentration of between 0.6 to 6 percent.
8. (original) The method of claim 1, wherein the detergent added is a pure detergent.
9. (original) The method of claim 1, wherein the host cells are at least 1 kg in weight.
- 10.(original) The method of claim 1, wherein the host cells are *Pichia pastoris* cells.
- 11.(original) The method of claim 1, wherein the buffer has an ionic strength of less than 350 mM and at least 1 mM.
- 12.(original) The method of claim 1, further comprising the step of incubating the cell and detergent mixture from 90 minutes to 24 hours.
- 13.(original) The method of claim 12, wherein the incubation is from 8 hours to 24 hours.
- 14.(original) The method of claim 1, further comprising the step of incubating the cell and detergent mixture at a temperature of between about 3°C and about 10°C.
- 15.(new) The method of claim 2, wherein the detergent is not dimethyltridecylamine.
- 16.(new) The method of claim 3, further comprising the step of adding glycerol to the suspended host cells, and wherein the glycerol is in a concentrated form of at least 3 times the desired final concentration of the glycerol after dilution in the buffer of step 1.

17.(new) The method of claim 4, wherein the detergent comprises a final concentration of between 0.1 to 10 percent.

18.(new) The method of claim 4, wherein the glycerol comprises a final concentration of between 0.6 to 15 percent.

19.(new) The method of claim 5, wherein the glycerol comprises a final concentration of between 0.6 to 6 percent.